arXiv:1611.04385v2 [math.NT] 17 Nov 2016

DOUGALL'S ${}_5F_4$ SUM AND THE WZ-ALGORITHM

JESÚS GUILLERA

Abstract. We show how to prove the examples of a paper by Chu and Zhang using the WZ-algorithm.

1. Introduction

Chu and Zhang had the good idea of accelerating the following Dougall's sum [1], which is convergent for $\mathrm{Re}(1+2a-b-c-d-e)>0$:

$$\sum_{k=0}^{\infty} \frac{(b)_k (c)_k (d)_k (e)_k}{(1+a-b)_k (1+a-c)_k (1+a-d)_k (1+a-e)_k} (a+2k),$$

using many recurrence relations patterns. Their paper has interesting formulas and many nice examples. We show that the Wilf-Zeilberger algorithm can do it automatically [2], and give four examples corresponding to Examples 42, 45, 50 and 53 of that paper. Note that Dougall's sum in [1] has two wrong signs and observe that it is symmetric in b, c, d and e (that is, these parameters are interchangeable).

We will use the following notation:

$$A(a,b,c,d,e,k) = \frac{(b)_k (c)_k (d)_k (e)_k}{(1+a-b)_k (1+a-c)_k (1+a-d)_k (1+a-e)_k} (a+2k).$$

Then Dougall's sum is

$$\Omega(a,b,c,d,e) = \sum_{k=0}^{\infty} A(a,b,c,d,e,k).$$

If $e \to -\infty$ then we have the sum

$$\Phi(a,b,c,d) = \sum_{k=0}^{\infty} B(a,b,c,d,k)$$

where

$$B(a,b,c,d,k) = (-1)^k \frac{(b)_k (c)_k (d)_k}{(1+a-b)_k (1+a-c)_k (1+a-d)_k} (a+2k).$$

This introduction is short and we have only cited two papers but the interested reader can also see the introduction and the references of [1].

2. Examples

In this section we prove the Examples 42, 45, 50 and 53 of the paper [1] using methods based on the WZ-algorithm, but we can derive all the other examples of that paper using the same kind of WZ-techniques.

Example 42. It accelerates

$$\Omega\left(\frac{3}{2},\frac{1}{2},1,1,1\right)=\frac{7}{4}\zeta(3),$$

see [1, Theorem 5], using the recurrence relation [3,0,1,2,2]. To do it with the WZ-method we consider the function

$$U(n,k)=A\left(\frac{3}{2}+3n,\frac{1}{2}+0n,1+1n,1+2n,1+2n,k\right),$$

and use the Maple package `with(SumTools[Hypergeometric])`. We see that the output of `Zeilberger(U(n,k),n,k,N)[1]` is equal to

$$-1024\left(\frac{1}{2}+n\right)(1+n)^5\,N+1728\left(\frac{3}{2}+n\right)^2\left(\frac{2}{3}+n\right)\left(\frac{4}{3}+n\right)\left(\frac{3}{4}+n\right)\left(\frac{5}{4}+n\right).$$

This output indicates that if

$$\begin{aligned}F(n,k)&=U(n,k)\frac{\left(\frac{1}{2}\right)_n(1)_n^5}{\left(\frac{3}{2}\right)_n^2\left(\frac{2}{3}\right)_n\left(\frac{4}{3}\right)_n\left(\frac{3}{4}\right)_n\left(\frac{5}{4}\right)_n}\left(-\frac{16}{27}\right)^n\\&=\frac{U(n,k)}{2n+1}\,\frac{(1)_n^5}{\left(\frac{3}{2}\right)_n\left(\frac{2}{3}\right)_n\left(\frac{4}{3}\right)_n\left(\frac{3}{4}\right)_n\left(\frac{5}{4}\right)_n}\left(-\frac{16}{27}\right)^n,\end{aligned}$$

then the output of `Zeilberger(F(n,k),n,k,N)[1]` is equal to $N-1$ (Observe that $-1024/1728=-16/27$). Hence, there is a function $G(n,k)$, such that (F,G) is a WZ-pair. Writing

`simplify(Zeilberger(F(n,k),n,k,N)[2]/F(n,k))`

we get a rational function $R(n,k)$ such that $G(n,k)=R(n,k)F(n,k)$. In addition, we see that

$$R(n,0)=\frac{172n^2+269n+106}{9(2+3n)(3+4n)},\quad U(n,0)=\frac{3}{2}+3n.$$

As the limit of $F(n,k)$ as $n\to\infty$ is equal to 0, we can use the Zeilberger's identity

$$\sum_{n=0}^{\infty}G(n,0)=\sum_{k=0}^{\infty}F(0,k)+\lim_{k\to\infty}\sum_{n=0}^{\infty}G(n,k).$$

We can easily check that in this example the limit commute with the summation and is equal to zero. Therefore, we have

$$\sum_{n=0}^{\infty}G(n,0)=\sum_{k=0}^{\infty}F(0,k)=\Omega\left(\frac{3}{2},\frac{1}{2},1,1,1\right)=\frac{7}{4}\zeta(3).$$

Hence

$$36\sum_{n=0}^{\infty}G(n,0)=\sum_{n=0}^{\infty}\frac{(1)_n^5}{\left(\frac{3}{2}\right)_n\left(\frac{5}{3}\right)_n\left(\frac{4}{3}\right)_n\left(\frac{7}{4}\right)_n\left(\frac{5}{4}\right)_n}\left(-\frac{16}{27}\right)^n(172n^2+269n+106)=63\zeta(3),$$

which is the formula of [1, Example 42].

Example 45. This example uses the same recurrence relation, that is [3,0,1,2,2]. We consider the function

$$U(n,k) = A\left(\frac{1}{2}+3n, \frac{1}{2}+0n, \frac{1}{2}+1n, \frac{1}{2}+2n, \frac{1}{2}+2n, k\right).$$

Then, procceeding as in the previous example, we obtain

$$F(n,k) = U(n,k)\cdot n\frac{\left(\frac{1}{2}\right)_n\left(\frac{1}{4}\right)_n^3\left(\frac{3}{4}\right)_n^3}{\left(\frac{1}{3}\right)_n\left(\frac{2}{3}\right)_n(1)_n^5}\left(-\frac{16}{27}\right)^n,$$

and the rational function $R(n,k)$ such that $G(n,k) = R(n,k)F(n,k)$. In addition, we get

$$R(n,0) = \frac{1376n^4+1808n^3+784n^2+138n+9}{16n(1+6n)(2+3n)(1+3n)}, \quad U(n,0) = \frac{1}{2}+3n.$$

We see that

$$\sum_{k=0}^{\infty} F(0,k) = \sum_{k=0}^{\infty}\left\{0\cdot A\left(\frac{1}{2},\frac{1}{2},\frac{1}{2},\frac{1}{2},\frac{1}{2},k\right)\right\} = 0, \quad \text{and} \quad \Omega\left(\frac{1}{2},\frac{1}{2},\frac{1}{2},\frac{1}{2},\frac{1}{2}\right) = \infty.$$

Therefore

$$\sum_{n=0}^{\infty} G(n,0) = \sum_{k=0}^{\infty} F(0,k) + \lim_{k\to\infty}\sum_{n=0}^{\infty} G(n,k) = \lim_{k\to\infty}\sum_{n=0}^{\infty} G(n,k).$$

We can check that the limit commutes with the summation, and in addition the limit of $G(n,k)$ as $k\to\infty$ is zero for $n\neq 0$. Hence

$$\sum_{n=0}^{\infty} G(n,0) = \lim_{k\to\infty} G(0,k) = \lim_{k\to\infty}\left(\frac{1}{32}\frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4}\frac{(2k+3)^2(2k+1)^3}{(k+1)^2(k+2)}\right) = \frac{1}{\pi^2}.$$

That is

$$\sum_{n=0}^{\infty}\frac{\left(\frac{1}{2}\right)_n\left(\frac{1}{4}\right)_n^3\left(\frac{3}{4}\right)_n^3}{\left(\frac{4}{3}\right)_n\left(\frac{5}{3}\right)_n(1)_n^5}\left(-\frac{16}{27}\right)^n(1376n^4+1808n^3+784n^2+138n+9) = \frac{64}{\pi^2},$$

which is the formula of [1, Example 45]. We will finally deduce that this example accelerates

$$9 - \frac{9}{16}\,\Omega\left(\frac{7}{2},\frac{1}{2},\frac{3}{2},\frac{5}{2},\frac{5}{2}\right).$$

For it observe that if $(F(n,k),G(n,k))$ is a WZ-pair then $(F(n+1,k),G(n+1,k)$ is also a WZ-pair. Hence

$$\sum_{n=0}^{\infty} G(n+1,0) = \sum_{k=0}^{\infty} F(1,k) + \lim_{k\to\infty}\sum_{n=0}^{\infty} G(n+1,k).$$

As the above limit is equal to zero, we obtain

$$\begin{aligned}\frac{1}{\pi^2} = \sum_{n=0}^{\infty} G(n,0) &= G(0,0) + \sum_{n=0}^{\infty} G(n+1,0) = G(0,0) + \sum_{k=0}^{\infty} F(1,k)\\ &= \frac{9}{64} - \frac{9}{1024}\,\Omega\left(\frac{7}{2},\frac{1}{2},\frac{3}{2},\frac{5}{2},\frac{5}{2}\right),\end{aligned}$$

and from it, we deduce that

$$\Omega\left(\frac{7}{2},\frac{1}{2},\frac{3}{2},\frac{5}{2},\frac{5}{2}\right) = 16 - \frac{1024}{9\pi^2},$$

which is a known result because Maple also gives the exact value of this Dougall's sum.

Example 50. It accelerates

$$\Phi\left(\frac{3}{2},1,1,1\right)=G \quad \text{(Catalan constant)},$$

see [1, Theorem 5], using the recurrence relation [3,0,2,2]. To do it with the WZ-method we consider the function

$$U(n,k)=B\left(\frac{3}{2}+3n,\frac{1}{2}+0n,1+2n,1+2n,k\right),$$

Then, with the Wilf-Zeilberger's algorithm, we obtain

$$F(n,k)=U(n,k)\frac{\left(\frac{1}{2}\right)_n^2(1)_n^2}{\left(\frac{3}{2}\right)_n^2\left(\frac{5}{6}\right)_n\left(\frac{7}{6}\right)_n}\left(\frac{16}{27}\right)^n=\frac{U(n,k)}{(2n+1)^2}\frac{(1)_n^2}{\left(\frac{5}{6}\right)_n\left(\frac{7}{6}\right)_n}\left(\frac{16}{27}\right)^n,$$

and the rational function $R(n,k)$ such that $G(n,k)=R(n,k)F(n,k)$. In addition, we get

$$R(n,0)=\frac{22n+21}{5+6n},\quad U(n,0)=\frac{1}{2}+3n,$$

and

$$\sum_{n=0}^{\infty}G(n,0)=\sum_{k=0}^{\infty}F(0,k)=\Phi\left(\frac{3}{2},1,2,2\right)=G.$$

Hence

$$30\sum_{n=0}^{\infty}G(n,0)=\sum_{n=0}^{\infty}\frac{(1)_n^2}{\left(\frac{7}{6}\right)_n\left(\frac{11}{6}\right)_n}\frac{22n+21}{2n+1}\left(\frac{16}{27}\right)^n=30\,G,$$

which is the formula in [1, Example 50].

Example 53. This example uses the recurrence relation [3,1,1,1,3]. From the function

$$U(n,k)=A\left(\frac{1}{2}+3n,\frac{1}{2}+n,\frac{1}{2}+n,\frac{1}{2}+n,3n,k\right),$$

we obtain

$$F(n,k)=U(n,k)\cdot n^2\frac{\left(\frac{1}{2}\right)_n^3\left(\frac{1}{3}\right)_n\left(\frac{2}{3}\right)_n}{(1)_n^5}\left(\frac{27}{64}\right)^n,$$

and the rational function $R(n,k)$ such that $G(n,k)=R(n,k)F(n,k)$. In addition, we get

$$R(n,0)=\frac{74n^2+27n+3}{16n^2(1+6n)},\quad U(n,0)=\frac{1}{2}+3n,$$

and

$$G(0,k)=\frac{3}{32}\frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2}\frac{(2k+1)^2}{(k+1)^2}.$$

Then, we use the Wilf-Zeilberger's identity

$$\sum_{n=0}^{\infty}G(n,k+1)=\sum_{n=0}^{\infty}G(n,k)+F(0,k)-\lim_{n\to\infty}F(n,k).$$

But $F(0,k)$ and the limit of $F(n,k)$ as $n\to\infty$ are equal to zero in this example. Therefore, we have

$$\sum_{n=0}^{\infty}G(n,k+1)=\sum_{n=0}^{\infty}G(n,k).$$

In addition, we can prove that Carlson's Theorem holds here. Hence, the summation above is not only periodic of period $k = 1$ but constant for every complex number k. That is

$$\sum_{n=0}^{\infty} G(n,k) = \text{constant} \quad \forall k \in \mathbb{C}.$$

To determine the constant, we write

$$\sum_{n=0}^{\infty} G(n,k) = \lim_{k\to -\frac{1}{2}} \sum_{n=0}^{\infty} G(n,k) = \lim_{k\to -\frac{1}{2}} G(0,k) = \frac{3}{2\pi^2},$$

because the limit commutes with the summation and is equal to 0 when $n \neq 0$. Hence

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^3 \left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n}{(1)_n^5} \left(\frac{27}{64}\right)^n (74n^2 + 27n + 3) = \frac{48}{\pi^2},$$

which is the formula [1, Example 53].

Remark. Some of the examples in [1], like the Example 45, have $1+2a-b-c-d-e \leq 0$, and then $\Omega(a,b,c,d,e)$ is not defined. Hence the left hand side of the formulas in the theorems of [1] has no sense in these cases and variants of the theorems would be needed. In this paper we have shown that the Wilf-Zeilberger's techniques can manage all the situations.

References

[1] W. Chu and W. Zhang Accelerating Dougall's ${}_5F_4$-sum and infinite series involving π, *Journal: Math. Comp.* **83** (2014), 475–512.

[2] M. Petkovšek, H. S. Wilf and D. Zeilberger, A=B, *A.K. Peters Ltd.* (1996).

Universidad de Zaragoza (Spain)
E-mail address: jguillera@gmail.com